




Differentially abundant KIRC/P miRNAs between GA2x and HiSeq

G. Robertson, Andy Chu, Richard Corbett
25 May 2011, 16h50

1. Adjusted data: GA2x vs. subsampled HiSeq.
2. Original data: GA2x vs. non-subsampled HiSeq
GC content of miRNAs






logConc	logFC	P.Value	adj.P.Val
hsa-mir-23c	-19.43	-2.20	8.01e-12 5.91e-09
hsa-mir-326	-14.64	0.991	2.39e-06 0.00088
hsa-mir-874	-15.61	0.961	9.25e-06 0.0023
hsa-mir-374c	-17.01	0.963	2.23e-05 0.00411
hsa-mir-1249	-19.48	1.30	4.74e-05 0.0070
hsa-mir-1976	-16.60	0.854	0.00014 0.017
...			




KIRC/P GA2x vs HiSeq, subsample 2, 735 miRNAs

edgeR v1.51






gene	logConc	logFC	P.Value	adj.P.Val
hsa-mir-23c	-19.40	-2.133	1.69e-11	1.24e-08
hsa-mir-326	-14.65	0.975	3.10e-06	0.00114
hsa-mir-874	-15.701	0.906	2.55e-05	0.00625
hsa-mir-374c	-17.033	0.911	5.31e-05	0.00813
hsa-mir-1249	-19.485	1.280	5.53e-05	0.00813
hsa-mir-1976	-16.611	0.797	0.00037	0.0450
...				






25 May 2011, 13h10

KIRC/P GA2x vs HiSeq, subsample 1, 735 miRNAs



gene	logConc	logFC	P.Value	adj.P.Val
hsa-mir-23c	-19.477	-2.29	2.88e-12	2.12e-09
hsa-mir-874	-15.620	0.984	5.62e-06	0.0021
hsa-mir-326	-14.679	0.923	1.13e-05	0.0024
hsa-mir-1249	-19.440	1.376	1.32e-05	0.0024
hsa-mir-374c	-17.047	0.884	0.000101	0.0148



KIRC/P GA2x vs HiSeq, subsample 1, 735 miRNAs



gene	logConc	logFC	P.Value	adj.P.Val
hsa-mir-23c	-19.477	-2.29	2.88e-12	2.12e-09
hsa-mir-874	-15.620	0.984	5.62e-06	0.0021
hsa-mir-326	-14.679	0.923	1.13e-05	0.0024
hsa-mir-1249	-19.440	1.376	1.32e-05	0.0024
hsa-mir-374c	-17.047	0.884	0.000101	0.0148

KIRC/P GA2x vs HiSeq, subsample 5, 35 miRNAs

Differential abundance analysis, edgeR v1.5.1





gene	logConc	logFC	P.Value	adj.P.Val
mi-13	-5.4255	0.783499	0.000945101	0.0330785
mi-23	-4.70622	0.406441	0.0291534	0.510185
mi-1	-3.89152	-0.221841	0.120596	0.751885
mi-2	-3.89345	-0.220216	0.120596	0.751885
mi-3	-3.87952	-0.220564	0.122239	0.751885
mi-34	-4.04362	0.211961	0.152072	0.751885
...				


1. Adjusted data: GA2x vs. subsampled HiSeq.
 2. Original data: GA2x vs. non-sampled HiSeq
- GC content of miRNAs

Differential abundance between HiSeq and GA2x

We identified differentially abundant miRNAs between HiSeq and GA2x (1/0) using edgeR v2.2.3 and an expression matrix of 767 miRNAs that had nonzero sums across 2x56 samples. Nine miRNAs had BH-corrected p-val < 0.05 (red in 'a', black squares in 'b', table). As expected, the FC plot ('smearPlot') identified no low-abundance miRNAs as differentially abundant. **hsa-mir-23c** was flagged as having a negative PC1 loading (slide 9), when all other 8 loadings had positive values.




gene	logConc	logFC	P.Value	adj.P.Val
→ hsa-mir-23c	-19.4337	-2.114	2.88E-11	2.21E-08
hsa-mir-326	-14.6473	1.072	1.52E-07	0.0000583
hsa-mir-874	-15.6407	1.031	1.01E-06	0.000257
hsa-mir-374c	-17.0304	1.000	6.31E-06	0.00121
hsa-mir-1249	-19.488	1.366	0.0000124	0.00190
hsa-mir-1976	-16.6154	0.8766	0.0000653	0.00833
hsa-mir-204	-11.4417	0.7045	0.000394	0.0431
hsa-mir-150	-11.2998	0.6859	0.000556	0.0481
hsa-mir-99b	-4.96516	0.6833	0.000565	0.0481



1. Open black circles/triangles show the 20 PCA 'driver' miRNAs. Squares show nine differentially abundant miRNAs for BH-pval<0.05.
2. Red and green point clouds appear to separate for mean abundances below ~2.

GC content of differentially abundant miRNA mature|star strands


GC content of 1733 mature or star strands (v17, blue) and 939 primary transcripts (v15, green)



miRNA	MIMAT	sequence	f(GC)
hsa-miR-874	MIMAT0004911	CUGCCCUGGCCCGAGGGACCGA	0.773
hsa-miR-326	MIMAT0000756	CCUCUGGGCCCUUCCUCCAG	0.700
hsa-miR-1976	MIMAT0009451	CCUCCUGCCCUCCUUGCUGU	0.650
hsa-miR-1249	MIMAT0005901	ACGCCCUUCCCCCCCUCUUCA	0.636
hsa-miR-99b	MIMAT0000689	CACCCGUAGAACCGACCUUGCG	0.636
hsa-miR-150	MIMAT0000451	UCUCCCAACCCUUGUACCAGUG	0.545
hsa-miR-23c	MIMAT0018000	AUCACAUUGCCAGUGAUUACCC	0.454
hsa-miR-204	MIMAT0000265	UUCCCUUUGUCAUCCUAUGCCU	0.454
hsa-miR-374c	MIMAT0018443	AUAAUACAACCUGCUAAGUGCU	0.364

/Users/grobertson/GENEREG/Data/miRNA/mirBase/v16/mature.hsa.fraction-GC.txt

Nine differentially abundant miRNAs (BH-pval<0.05, red) for 767 miRNAs (blue) in GA2x-HiSeq KIRC/KIRP test



gene	logConc	logFC	P.Value	adj.P.Val
hsa-mir-23c	-19.43	-2.114	2.88E-11	2.21E-08
hsa-mir-326	-14.64	1.072	1.52E-07	0.0000583
hsa-mir-874	-15.640	1.031	1.01E-06	0.000257
hsa-mir-374c	-17.030	1.000	6.31E-06	0.00121
hsa-mir-1249	-19.488	1.366	0.0000124	0.00190
hsa-mir-1976	-16.615	0.877	0.0000653	0.00833
hsa-mir-204	-11.442	0.705	0.000394	0.0431
hsa-mir-150	-11.30	0.686	0.000556	0.0481
hsa-mir-99b	-4.965	0.683	0.000565	0.0481